
Management Plan for the improvement of Bobwhite Quail (Colinus virginianus) Habitat

Aliyah Abdur-Razzaaq




Introduction

- Northern Bobwhite is a species of quail native to the United States.
- Over the last 35 years in bobwhite populations have declined by more than 75% in Texas
- Potential causes of decline: drought, land use changes, land fragmentation, habitat loss, invasive species, insecticides, diseases and parasites.



Habitat/Lifestyle

- Agricultural fields, grasslands, pine forests, and grass-brush rangelands.
- Open ground cover ~30% to 50% vegetative coverage.
- Avoid mature woodlands and prefer the early stages of woody regrowth after a fire or other disturbance
- Prefer places with access to water
- Ground Nesters
 - Mainly nest in bunchgrasses
- Winter/Fall Diet mainly seeds
 - Forbs and grasses
- Spring/Summer Diet mainly insects and plant mast



(Ivanov, Northern Bobwhite Quail)

Objectives

To be implemented after comparison between Savanna C North and Savanna D South. The location most suitable for the following objectives will be chosen.

- 1. Monitoring the population of little bluestem once per year to ensure frequency does not decrease by more than 25%.
- 2. Ingrease bare ground cover from current measurement to 30%
 - a. To be implemented via controlled burns and cattle grazing
 - b. Sampling for cover of bare ground to occur once per year

Site Descriptions

Savanna C North

- Located in Ecoregion 3 Post oak Savannah.
- Open grassland containing a small pond.
 Soil is majority composed of Boonville fine sandy loam (70+%)
- High Runoff class
- Last burn treatment march 2018

Savanna D South

- Located in Ecoregion 3 Post oak Savannah .
- Forested area containing a small pond.
- Soil composed of Zack fine sandy loam (100%)
- High Runoff Class
- Last burn treatment April 2018

Methods: Sampling

Savanna C North

Frequency

- Mixture of Line Intercept and Random Quadrats (60 total)
- Total hits measure
- ,25m² frame used

Cover

- Random Quadrats
- Percent cover measured
- .25m² frame used

Savanna D South

Frequency

- Mixture of Line Intercept and Random Quadrats (60 total)
- Total hits measured
- .25m² frame used

Cover

- Point Intercept
- Number of hits measured
- Stake/Flag used

Methods: Analysis

- Each method is compared with an error rate of .05%
- Frequency of little bluestem Savanna C North and Savanna D south compared using chi-square analysis.
- Frequency of trees/shurbs in Savanna C North and Savanna D south compared using chi-square analysis.
- Cover unable to be compared directly due to different sampling methods.

 Instead:
 - Cover in Savanna C North compared to the ideal mean using a single sample T-test
 - Cover in Savanna D south compared to the ideal number of hits using chi-square analysis.

Results

Little bluestem chi-square

- Null Hypothesis: Both plots of land contain the same amount of little bluestem
- Alternative: Savanna D south contains more little bluestem than Savanna C north

Critical value: 3.84

Significance level: .05

P value: . 007244

Null Hypothesis rejected

Savanna C North

.			
name	hits		percent
ANTR	11113	2	3.33%
BULA		1	1.67%
		1	1.67%
CAIN		1	
CYPER			1.67%
DIOL		4	6.67%
ERCA		1	1.67%
EUPAT		5	8.33%
ILVO		5	8.33%
LALIPO		1	1.67%
NALE		3	5.00%
NOST		1	1.67%
PAPL		2	3.33%
RUTR		13	21.67%
SCSC		47	78.33%
SMILA		1	1.67%
SPCR		1	1.67%
ULCR		1	1.67%
ULMUS		1	1.67%
VERBI		3	5.00%
VISCI		1	1.67%
YUCCA		1	1.67%
TOTAL		47	

Savanna D south

<u> </u>		J	30111
	Number o		
name	hits		percent
ANTE		1	1.67%
BAPTI		3	
BOTHR		2	3.33%
CLOVER		1	1.67%
CRTE		1	1.67%
ILVO		1	1.67%
JUVI		1	1.67%
LUTE		1	1.67%
NALE		6	10.00%
NOBI	1	1	18.33%
OPUNT		1	1.67%
PAFL		2	3.33%
QUST		2	3.33%
RUTR		2	3.33%
SCSC	5	7	95.00%
TOTAL	57		
IOIAL	<i>31</i>		

Results con't

- Tree/Forb chi-square
- Null Hypothesis: Both plots of land contain the same amount of Trees & shrubs
- Alternative: Savanna C North contains Trees/Shrubs than Savanna D south

Critical value: 3.84

Significance level: .05

P value: .004485

Null Hypothesis rejected

Savanna C North

.			•
namo	hits		percent
name ANTR	11115	2	3.33%
BULA		1	1.67%
		1	1.67%
CAIN		1	1.67%
CYPER		4	6.67%
DIOL		1	1.67%
ERCA			
EUPAT		5	8.33%
ILVO		5	8.33%
LALIPO		1	1.67%
NALE		3	5.00%
NOST		1	1.67%
PAPL		2	3.33%
RUTR		13	21.67%
SCSC		47	78.33%
SMILA		1	1.67%
SPCR		1	1.67%
ULCR		1	1.67%
ULMUS		1	1.67%
VERBI		3	5.00%
VISCI		1	1.67%
YUCCA		1	1.67%
TOTAL		20	

Savanna D south

Javan		50111
	Number of	
name		percent
ANTE	1	1.67%
BAPTI	3	5.00%
BOTHR	2	3.33%
CLOVER	1	1.67%
CRTE	1	1.67%
ILVO	1	1.67%
JUVI	1	1.67%
LUTE	1	1.67%
NALE	6	10.00%
NOBI	11	18.33%
OPUNT	1	1.67%
PAFL	2	3.33%
QUST	2	<mark>3.33%</mark>
RUTR	2	<mark>3.33%</mark>
SCSC	57	95.00%
TOTAL	7	
101/12	•	

Cover chi-square

		Total
Cover	Amount	Amount
Classification	Counted	Possible
Bare Ground	9	
Litter	5	
Grass	12	
Forb	1	
Woody	2	29
IDEAL bare		
ground	15	

- Null Hypothesis: Savanna D south has the same amount of bare ground cover as the Ideal.
- Alternative: The ideal contains more bare ground cover than Savanna D south.

Critical value: 3.84

Significance level: .05

P value: .109682

Fail to Reject Null Hypothesis

Cover single sample T-test

Mean	4.3
standard deviation	6.447221624
Variance	41.56666667
Observations	10
SEM	2.038790491
hypothesized	
mean (ideal)	50
df	9

- Null Hypothesis: Savanna C North has the same amount of bare ground cover as the Ideal.
- Alternative: The ideal contains more bare ground cover than Savanna C North.

Alpha level 0.05 t Stat -22.41525071

t dist 0.00000003314605257

Null Hypothesis Rejected

Implications & Actions

- Savanna C North
- Contains less little bluestem
- Contains more trees/shrubs
- Contains less bare ground than the Ideal.

- Savanna D South
- Contains more little
 Bluestem
- Contains less tree/shrubs
- Same amount of Bare ground as Ideal

For monitoring purposes Savanna D south would be more ideal to implement controlled burn and management of little bluestem than Savanna C North. Starting with a baseline of more little bluestem will make it easier to prevent a decline of over 25%. However, it is important to note point intercept is not the best metric to measure cover. This may have influenced the results.

Budget & Timeline

- Estimated timeline approx.~ 5 years.
- Budgeted to reflect 4 people monitoring for 2 weeks(1 week before burn and 1 week after 6 months)
- Cattle Costs accounts for 1 week of grazing.

	Cost
Controlled burn	1914\$
Personell	800\$
Equipment	2000\$
Monitoring	38,400
cattle	2380\$
Total	45,494\$

Literature Cited

- Birds of the Oaks & Prairies and Osage Plains of Texas. https://www.cedarhillstatepark.org/nature/animals/oakprariebirdlist.pdf.
- Bobwhite Quail Management, https://tpwd.texas.gov/landwater/land/habitats/high_plains/upland_game/bobwhite.pht ml.
- Brown, Sarah K, and William E. Palmer. "Effect of Prescribed Fire and Mechanical Treatments on Northern Bobwhite Occupancy in Mesic Pine Flatwoods." National Quail Symposium Proceedings, vol. 9, 2022, https://doi.org/10.7290/nqsp09vnfj.
- Couvillon, Ross O., et al. "Habitat Space Used by Northern Bobwhites and Texas Tortoises on South Texas Rangelands." Quail, vol. 8, 2017, https://doi.org/10.7290/nqsp08w49n.
- Hernandez, Fidel, et al. "The Forgotten Quail Decline: The Plight of Scaled Quail in Texas." Quail, vol. 7, 2017, https://doi.org/10.7290/nqsp07h255.
- Johnston, Bradley K., et al. "Bobwhite Response to Cattle Grazing in South Texas." National Quail Symposium Proceedings, https://doi.org/10.7290/nqsp.
- "Potential Causes of the Texas Quail Decline." Potential Causes of the Texas Quail Decline | Texas A&M NRI, https://nri.tamu.edu/publications/extension-publications/2018/potential-causes-of-the-texas-quail-decline/.
- "Texas Ecoregions." Texas Ecoregions Texas Parks & Wildlife Department, 16 Dec. 2022, https://tpwd.texas.gov/education/hunter-education/online-course/wildlife-conservation/texas-ecoregions.